

Susceptibility of House Fly (*Diptera: Muscidae*) Larvae to Entomopathogenic Nematodes (*Rhabditida: Heterorhabditidae, Steinernematidae*)

DAVID B. TAYLOR, ALLEN L. SZALANSKI,¹ BYRON J. ADAMS,¹ AND RICHARD D. PETERSON II

Midwest Livestock Insects Research Laboratory, USDA-ARS, Department of Entomology, University of Nebraska, Lincoln, NE 68583

Environ. Entomol. 27(6): 1514-1519 (1998)

ABSTRACT The potential for entomopathogenic nematodes to control flies in cattle feedlots was determined by screening 40 strains representing 8 species of *Heterorhabditis* Poinar and 5 species of *Steinernema* Travassos for virulence toward 3rd-instar house flies (maggots), *Musca domestica* L. None of the 22 strains of *Heterorhabditis* infecting maggots caused significant levels of mortality in a filter paper assay. Ten strains of *Steinernema* infected maggots, of which 7 strains (4 *S. carpocapsae* (Weiser), 2 *S. feltiae* (Filipjev), and 1 *S. scapterisci* Nguyen & Smart) caused significant mortality. Ten *Heterorhabditis* strains and 10 *Steinernema* strains successfully reproduced for ≥ 2 generations in maggots. No difference was observed between 72-h survival of maggots and adult emergence. Six strains of *Steinernema* were selected for 10 generations on maggots and then compared with unselected lines. No difference in pathogenicity between selected and unselected lines was observed. Two strains of *S. feltiae*, SN and UNK-36, and 2 of the best *Heterorhabditis* strains, *H. bacteriophora* Poinar OSWEGO and *H. megidis* Poinar, Jackson & Klein HF-85 were tested in a fresh bovine manure substrate. All 4 strains produced significant fly mortality in the manure substrate, although the *S. feltiae* strains had significantly lower LC_{50} values than did the *Heterorhabditis* spp. The most promising strain, *S. feltiae* SN, gave LC_{50} and LC_{99} values of 4 and 82 infective juveniles per maggot, respectively. These doses were equivalent to 2.7 and 55 infective juveniles per gram of manure and 5.1 and 104 infective juveniles per square centimeter of surface area. Infective juveniles capable of infecting greater wax moth larvae, *Galleria mellonella* (L.), survived in manure for up to 10 wk without hosts.

KEY WORDS *Musca domestica*, biological control, feedlot, entomopathogenic nematodes

NEMATODES OF THE genera *Heterorhabditis* Poinar and *Steinernema* Travassos, in conjunction with their symbiotic bacteria, *Photorhabdus* spp. and *Xenorhabdus* spp., are pathogenic to many insect pests (Poinar 1979) and are effective biological control agents for soil associated insects (Klein 1990). Several studies have addressed the efficacy of entomopathogenic nematodes against the house fly, *Musca domestica* L., in the laboratory and confined poultry environment (Renn et al. 1985; Geden et al. 1986; Belton et al. 1987; Mullens et al. 1987a, b). Although moderate levels of mortality were observed in petri dish-filter paper assays, virulence was greatly reduced when house fly larvae (maggots) were exposed in chicken manure substrates. Georgis et al. (1987) demonstrated that poultry manure was toxic to infective nematode larvae.

The filth fly breeding substrate in the cattle feedlot environment differs significantly from that of the confined poultry environment. Nitrogenous wastes are

excreted by cattle as water soluble urea, whereas they are in the form of uric acid in poultry manure. Also, the manure pit of a poultry house would tend to be relatively free of plant material and soil. Manure in feedlots is frequently mixed with soil, grain, and haylage. These differences may make the filth fly substrate in the cattle feedlot environment more hospitable to entomopathogenic nematodes than that in confined poultry systems. Despite these differences, the efficacy of entomopathogenic nematodes for the control of flies in the confined cattle environment has not been explored (Petersen 1993). In addition, few of the available species and strains of nematodes have been screened against filth flies, and conflicting results have been obtained for some of the nematode species, possibly because of differences in strain pathogenicity (Laumond et al. 1979, Gaugler 1988).

The potential for entomopathogenic nematodes to control filth flies in cattle feedlots was determined by screening 40 strains representing 8 species of *Heterorhabditis* and 5 species of *Steinernema* for pathogenicity toward maggots. Strains that successfully reproduced in maggots were reared for 10 generations on maggots in an effort to increase pathogenicity toward this host. Finally, the most promising strains were

This article reports the results of research only. Mention of a proprietary product does not constitute an endorsement or a recommendation by USDA for its use.

¹ Department of Plant Pathology, University of Nebraska, Lincoln, NE 68583

evaluated for pathogenicity and persistence in bovine manure.

Materials and Methods

Nematode and Fly Cultures. Species, strains, and sources of nematodes used are presented in Table 1. Classification of nematode strains to species follows that of Poinar (1990). Nematodes were reared using late-instar greater wax moths, *Galleria mellonella* (L.), obtained from Grubco (Hamilton, OH) using methods similar to Dutky et al. (1964). Nematodes were harvested with White traps (White 1927), quantified by counting the number of nematodes in five, 5- μ l samples and using the average to dilute to solutions to the needed concentration, and stored in tissue culture flasks at 15°C.

House flies were from a 9-yr-old colony collected from a feedlot in the vicinity of Lincoln, NE. Fly larvae were reared on a wheat bran diet at 25°C. Third instars were obtained for all experiments by removing them from the medium after 3 d of development.

Susceptibility of Maggots on Filter Paper. The virulence of *Heterorhabditis* spp. and *Steinernema* spp. toward maggots was tested using a filter paper assay. Fifteen 3rd instars were placed in a 9-cm petri dish lined with a filter paper disk (7 cm diameter). One milliliter of nematode-water suspension (\approx 100 nematodes per maggot) was applied to the filter paper. Water was used for control replicates. Maggot mortality was recorded at 3 d and adult fly emergence was scored at 2 wk. Reproductive success of the nematodes was evaluated by placing cadavers in White traps 7 d after infection. White traps were maintained at 25°C and checked daily for the emergence of infective juvenile nematodes. Nematodes were collected from the White traps and stored in tissue culture flasks at 15°C. Nematodes remained viable for >1 yr under these conditions, but cultures for bioassays were all <1 mo old.

Selection. Strains that successfully reproduced in maggots were maintained on that host for as many generations as possible. General procedures were the same as outlined in the susceptibility test except 25 maggots were used in each petri dish. Infections were attempted 3 times per generation. If, after 3 attempts, the nematodes failed to reproduce, that strain was dropped from the selection study. After selection for 10 generations, selected lines were compared with lines of the same strain maintained in greater wax moth larvae. The concentrations of nematodes used for this comparison were 500, 125, and 15 nematodes per maggot. Otherwise, procedures were the same as for the susceptibility study. To reduce any effect that parental host might have on pathogenicity, progeny of the original comparisons were tested for 2 additional generations giving comparisons of F_{10} with F_0 , F_{11} with F_1 , and F_{12} with F_2 selected lines.

Manure Substrate. Two strains of *Steinernema feltiae* (Filipjev), *H. bacteriophora* OSWEGO, and *H. megidis* HF-85, were tested for pathogenicity toward maggots in fresh bovine manure. Manure (<48 h old)

was collected from the ground. Manure was placed in a pan and mixed until uniform. Samples of manure (150 g wet weight) were partitioned into plastic containers (11 cm diameter, 8 cm deep) with screened holes (7.5 cm diameter) in the lid. Two replicates of 100 three-d-old maggots per strain were inoculated with 15,500, 7,500, 3,500, 1,500, 700, or 300 nematodes per container. The different nematode densities were run in successive weeks. The 2 *Heterorhabditis* spp. were tested at the highest 2 doses, *S. feltiae* UNK-36 was tested at the highest 4 doses and *S. feltiae* SN was tested at all 6 doses. Two weeks after initiation, house fly adults were counted in each container. Persistence of nematodes in the manure substrate was examined by placing 5 greater wax moth larvae in each container for 48 h each wk beginning 3 wk after initiation for the 15,500 and 7,500 tests. After 48 h, wax moth larvae were removed to 9-cm petri dishes lined with a piece of moistened filter paper (7 cm diameter) and scored for mortality 2 d later. Dead wax moth larvae were observed for nematode infections. Wax moth larvae from which no nematodes emerged were dissected to check for the presence of nematodes. Water was added to the manure containers once per week to maintain their original weight.

Data Analysis. Filter paper assays were conducted for 3 successive weeks with 3 replications of each nematode strain per week for a total 9 replicates of 15 maggots each per strain. A 1-way analysis of variance (ANOVA) with a complete block design (PROC GLM, SAS Institute, 1985) was used to analyze maggot mortality and adult emergence. Mortality data are presented as percentage mortality, although actual mortality was used for statistical tests. A *t*-test was used to separate the means, and Dunnett *t*-test (PROC GLM, SAS Institute 1985) was used to compare treatments with controls.

Probit analysis (PROC PROBIT, SAS Institute 1985) was used to calculate LC_{50} values for the selection and manure tests using a natural log transformation of the nematode concentrations. Overlap of the 95% fiducial limits was used to determine significance (Fuxa 1987).

Results

Seven *Steinernema* strains—*S. feltiae* SN and UNK-36, *S. carpocapsae* UNK-34, Kapow, Mex-1 and Mex-2, and *S. scapterisci*—produced significant mortality of maggots in the filter paper assay (Table 1, $F = 20.37$; $df = 42, 326$; $P \leq 0.001$). An additional 3 strains of *Steinernema* and 22 strains of *Heterorhabditis* produced low levels of infection, but mortality did not differ significantly from the control. Five *Steinernema* and 5 *Heterorhabditis* strains failed to infect maggots. Maggot survival and adult emergence did not differ significantly ($F = 1.99$; $df = 1, 78$; $P = 0.162$). Very few of the maggots alive after 3 d failed to pupate and emerge as adults. Dissections of unemerged pupae revealed no nematodes. Control mortalities were 0% for the 3-d maggot count and averaged 3% for the adult emergence count. Inspection of cadavers for several weeks following infection revealed successful repro-

Table 1. Species and strains of *Heterorhabditis* spp. and *Steinernema* spp. evaluated for virulence to house fly larvae

Species	Strain	Origin	Source	% mortality \pm SD at 72 h	Group	Reproduced
A. <i>Heterorhabditis</i> spp.						
<i>H. argentinensis</i> Stock	RAF	Rafaela, Argentina	Stock ^a	5.2 \pm 5.5	D	—
<i>H. argentinensis</i> Stock	JJ	Rabuela, Argentina	Jackson ^b	0.0 \pm 0.0	D	—
<i>H. bacteriophora</i> Poinar	NC-1	North Carolina	Brooks ^c	0.7 \pm 2.2	D	+
<i>H. bacteriophora</i> Poinar	OH-25	Oregon	Liu ^d	1.5 \pm 2.9	D	+
<i>H. bacteriophora</i> Poinar	OH-23	Oregon	Liu ^d	3.7 \pm 5.9	D	+
<i>H. bacteriophora</i> Poinar	lewiston	North Carolina	Jackson ^b	0.7 \pm 2.2	D	+
<i>H. bacteriophora</i> Poinar		Brecon, Australia	Akhurst ^e	0.0 \pm 0.0	D	—
<i>H. bacteriophora</i> Poinar	AB Road	Nebraska	Adams ^f	2.9 \pm 4.9	D	+
<i>H. bacteriophora</i> Poinar	A cows	Nebraska	Adams ^f	2.2 \pm 3.3	D	+
<i>H. bacteriophora</i> Poinar	OSWEGO	North Carolina	Jackson ^b	4.4 \pm 5.8	D	+
<i>H. bacteriophora</i> Poinar	Riwaka	New Zealand	Burnell/Joyce ^g	4.5 \pm 2.9	D	—
<i>H. bacteriophora</i> Poinar	HP88	Utah	Westerman ^h	0.0 \pm 0.0	D	—
<i>H. hawaiiensis</i> Gardner, Stock & Kaya	MG3 prime	Hawaii	Stock ^a	1.5 \pm 2.9	D	—
<i>H. hepialus</i> Stock, Strong & Gardner	—	California	Stock ^a	5.2 \pm 6.5	D	—
<i>H. indicus</i> Poinar, Karunakar & Hastings	—	Coimbatore, India	Burnell/Joyce ^g	0.0 \pm 0.0	D	—
<i>H. marelatus</i> Liu & Berry	OH-10	Oregon	Liu ^d	5.9 \pm 7.0	D	—
<i>H. megidis</i> Poinar, Jackson & Klein	HO-1	Ohio	Akhurst ^e	4.4 \pm 3.3	D	—
<i>H. megidis</i> Poinar, Jackson & Klein	Y	Netherlands	Smits ⁱ	0.7 \pm 2.2	D	—
<i>H. megidis</i> Poinar, Jackson & Klein	HF-85	Netherlands	Westerman ^h	3.7 \pm 8.9	D	+
<i>H. megidis</i> Poinar, Jackson & Klein	NLH94.1	Netherlands	Smits ⁱ	4.5 \pm 5.8	D	+
<i>H. megidis</i> Poinar, Jackson & Klein	X	British Columbia, Canada	Smits ⁱ	1.5 \pm 2.9	D	—
<i>H. megidis</i> Poinar, Jackson & Klein	NLHE87.3	Netherlands	Smits ⁱ	2.9 \pm 4.9	D	—
<i>H. megidis</i> Poinar, Jackson & Klein	Sp-Y	Europe	Smits ⁱ	0.7 \pm 2.2	D	—
<i>H. zealandica</i> Poinar	NZH	New Zealand	Akhurst ^e	4.4 \pm 3.3	D	+
<i>H. sp.</i>	OH-31	Oregon	Liu ^d	0.0 \pm 0.0	D	—
<i>H. sp.</i>	K122	Ireland	Griffin ^j	2.9 \pm 4.9	D	—
<i>H. sp.</i>	OH-1	Oregon	Liu ^d	2.2 \pm 3.3	D	—
B. <i>Steinernema</i> spp.						
<i>S. carpocapsae</i> (Weiser)	UNK-34	South Dakota	Jackson ^b	42.9 \pm 27.9	AB	+
<i>S. carpocapsae</i> (Weiser)	Mex-1	Mexico	Jackson ^b	15.5 \pm 17.3	C	+
<i>S. carpocapsae</i> (Weiser)	Mex-2	Mexico	Jackson ^b	17.8 \pm 4.7	C	+
<i>S. carpocapsae</i> (Weiser)	Agriotos	Aeriobo, Spain	Jackson ^b	1.5 \pm 2.9	D	+
<i>S. carpocapsae</i> (Weiser)	Kapow	Kapow, Poland	Jackson ^b	36.3 \pm 28.3	B	+
<i>S. carpocapsae</i> (Weiser)	UK	United Kingdom	Jackson ^b	5.2 \pm 6.5	D	+
<i>S. feltiae</i> (Filipjev)	UNK-36	South Dakota	Jackson ^b	49.6 \pm 25.4	A	+
<i>S. feltiae</i> (Filipjev)	SN	France	Jackson ^b	50.4 \pm 13.8	A	+
<i>S. glaseri</i> (Steiner)	—	—	Jackson ^b	0.0 \pm 0.0	D	—
<i>S. riobravisi</i> Cabanillas, Poinar & Raulston	—	Texas	Cabanillas ^k	0.0 \pm 0.0	D	—
<i>S. scapterisci</i> Nguyen & Smart	—	—	Jackson ^b	35.5 \pm 27.5	B	+
<i>S. sp.</i>	ARP blowout	Nebraska	Adams ^f	0.0 \pm 0.0	D	+
<i>S. sp.</i>	ARAP YS7	Nebraska	Adams ^f	0.0 \pm 0.0	D	—
C. Control	—	—	—	0.0 \pm 0.0	D	—

Data are the mean percentage of house fly larvae killed in filter paper assays. Each species was replicated 9 times with 15 house fly larvae per replicate. Groupings are based upon *t*-tests using least significant difference.

^a Stock, S. P., University La Plata CTR Parasities and Vectors Studies, La Plata, Argentina.

^b Jackson, J. J., Northern Grain Insects Research Laboratory, USDA-ARS, Brookings, SD.

^c Brooks, M. A., Department of Entomology, University of Minnesota, St. Paul, MN.

^d Liu, J., Department of Entomology, Oregon State University, Corvallis, OR.

^e Akhurst, R. J., CSIRO Division of Entomology, GPO Box 1700 Canberra, ACT 2601 Australia.

^f B.J.A.

^g Burnell, A. M., and S. A. Joyce, Department of Biology, St. Patricks College, Maynooth, Kildare, Ireland.

^h Westerman, P. R., Agrar. Hogeschool Friesland, P.O. Box 1528 8901 BV Leeuwarden, Netherlands.

ⁱ Smits, P. H. Research Institute Plant Protection, DLO, POB 9060, Wageningen NL-6700 GW, Netherlands.

^j Griffin, C. T., Department of Biology, St. Patricks College, Maynooth, Kildare, Ireland.

^k Cabanillas, H. E. Crop Insects Research Laboratory, Subtropical Agriculture Research Laboratory, USDA-ARS, Weslaco, TX.

duction and new infective juveniles for 20 nematode strains—10 *Steinernema* and 10 *Heterorhabditis* (Table 1).

Six strains of *Steinernema* reproduced for 10 generations in maggots (Table 2). Comparison of selected and unselected lines indicated that the selection procedure had no effect on the virulence of the nematodes toward house fly. LC₅₀ values for *S. feltiae* ranged from 10 to 18 infective nematodes per host, whereas those for *S. carpocapsae* and *S. scapterisci*

ranged between 38 and 55 infective nematodes per host.

Two strains of *S. feltiae* and 2 *Heterorhabditis* species (Table 3) were evaluated for their virulence toward maggots in manure. Each caused a reduction in the number of house fly adults emerging, although LC₅₀ values for the *Heterorhabditis* species were higher than those for *S. feltiae*. The LC₅₀ for *S. feltiae* UNK-36 in manure (30 infective juveniles per maggot) did not differ from that observed in the filter paper assay.

Table 2. Pathogenicity of selected and unselected *Steinernema* strains toward house fly larvae

Species	Strain	Selected	No. hosts	Slope ± SE	LC ₅₀ ^a	95% CL	χ ²
<i>S. feltiae</i>	UNK-36	-	389	0.64 ± 0.07	18	13-24	95.16
		+	401	0.68 ± 0.07	14	10-19	83.35
<i>S. feltiae</i>	SN	-	397	0.60 ± 0.07	10	6-14	65.51
		+	391	0.85 ± 0.11	11	8-15	52.53
<i>S. carpocapsae</i>	UNK-34	-	404	0.59 ± 0.06	30	22-39	111.49
		+	396	0.62 ± 0.06	28	21-36	113.30
<i>S. carpocapsae</i>	Mex	-	391	0.51 ± 0.10	55	44-69	26.06
		+	403	0.69 ± 0.07	40	31-50	138.88
<i>S. carpocapsae</i>	Kapow	-	400	0.57 ± 0.05	47	37-58	156.32
		+	402	0.84 ± 0.07	33	26-40	143.54
<i>S. scapterisci</i>		-	399	0.53 ± 0.05	43	31-56	107.50
		+	394	0.65 ± 0.06	37	28-47	132.50

^a Number of nematodes per host.

However, the LC₅₀ for *S. feltiae* SN in manure (4 infective juveniles per maggot) was about one-half the value observed on filter paper. LC₉₉ and LC₉₅ values for *S. feltiae* SN were 82 and 58 nematodes per host or 104 and 74 nematodes per square centimeter of manure surface area, respectively.

Survival of infective stage larval nematodes in manure was examined by placing 5 greater wax moth larvae in cups inoculated with 15,500 or 7,500 and 100 maggots beginning 3 wk after inoculation. No hosts were available for nematode reproduction after the initial 100 maggots. Infective juveniles of the *Heterorhabditis bacteriophora* and *S. feltiae* persisted for 10 wk (Table 4). Mortality of the wax moth larvae varied greatly, even within treatments. Therefore, the mortality data were not subjected to statistical analysis.

Discussion

Evaluation of 40 strains of entomopathogenic *Heterorhabditis* and *Steinernema* nematodes revealed 7 strains of *Steinernema* spp. that produced significant levels of house fly mortality (Table 1). *S. carpocapsae*, *S. feltiae*, and *S. scapterisci* were the most promising; however, strains of *S. carpocapsae* varied greatly in their virulence. Of the 6 strains of *S. carpocapsae* eval-

uated, 2 produced relatively high mortality, 2 moderate mortality, and 2 very low mortality. Both strains of *S. feltiae* were highly effective as was the 1 strain of *S. scapterisci*. None of the *Heterorhabditis* spp. showed promise for use in house fly control, although several strains successfully reproduced.

No differences were observed between maggot survival and adult emergence. This facilitated the later tests to evaluate house fly susceptibility to nematodes in manure substrates. Isolating maggots from these media can be tedious. Adult emergence gave us a more efficient method to measure nematode induced mortality.

Care must be used when comparing these results with those of previous studies at the species level because of changes in taxonomy. *Steinernema feltiae* of previous authors (Renn et al. 1985, Geden et al. 1986, Mullens et al. 1987a) were actually *S. carpocapsae*, whereas *S. bibionis* of those authors were *S. feltiae* (Poinar 1990). Among the *Heterorhabditis* spp., *H. heliothidis* of Renn et al. (1985) were *H. zealandica*, and *H. heliothidis* of Geden et al. (1986) and Mullens et al. (1987a) were *H. bacteriophora*.

Steinernema feltiae was the most virulent species toward maggots followed by *S. carpocapsae* in the filter paper assay. This differs from that of Mullens et al.

Table 3. Number of adult house flies emerging from bovine manure substrate inoculated with 100 house fly larvae and 155-3 infective *Heterorhabditis* spp. or *Steinernema feltiae* per fly larva

Species	Strain	No. infective nematodes per fly larva					Probit analysis			
		155 ^a	75 ^b	35 ^c	15 ^d	7 ^e	3 ^f	LC ₅₀ ^g	95% CL	Slope
<i>H. bacteriophora</i>	OSWEGO	4.0 ± 1.0*	60.5 ± 0.5*					83	78-89	1.754 ± 0.583
<i>H. megidis</i>	HF-85	10.5 ± 0.5*	81.0 ± 1.0*					106	101-113	2.127 ± 0.770
<i>S. feltiae</i>	UNK-36	0.0 ± 0.0*	1.0 ± 1.0*	51.5 ± 2.5*	68.5 ± 6.5			30	5-63	1.372 ± 0.450
<i>S. feltiae</i>	SN	0.0 ± 0.0*	0.0 ± 0.0*	20.0 ± 3.4*	22.7 ± 15.5*	33.5 ± 2.5*	61.5 ± 3.5*	4	3-5	0.776 ± 0.051
Control	-	75.5 ± 12.5	96.5 ± 2.5	89.5 ± 3.5 ^h		94.5 ± 1.5 ⁱ		-	-	-

* Differs significantly at P < 0.05 from control, Dunnett t-test.

^a F = 33.4; df = 4, 5; P = 0.001.

^b F = 1,192.7; df = 4, 5; P < 0.001.

^c F = 171.2; df = 2, 3; P = 0.001.

^d F = 36.2; df = 2, 3; P = 0.008.

^e F = 169.5; df = 1, 2; P = 0.006.

^f F = 75.1; df = 1, 2; P = 0.013.

^g No. infective juvenile nematodes per host.

^h One control was used for the 35 and 15 nematodes per maggot treatments.

ⁱ One control was used for the 7 and 3 nematodes per maggot treatments.

Table 4. Mean number of greater wax moth larvae dead after 48-h exposure to manure after initial inoculation with 100 house fly larvae and 155 or 75 *Heterorhabditis* spp. or *S. feltiae* per fly larva

Species	Strain	Weeks after inoculation							
		3	4	5	6	7	8	9	10
<i>H. bacteriophora</i>	OSWEGO	3.75 ± 1.64 ^a	1.50 ± 1.12 ^a	0.00 ± 0.00	1.50 ± 1.50 ^a	1.25 ± 1.09 ^a	1.25 ± 1.30	0.75 ± 0.83	1.00 ± 1.00 ^a
<i>H. megidis</i>	HF-85	3.50 ± 1.66 ^a	2.75 ± 1.48 ^a	0.25 ± 0.43	0.75 ± 0.83	1.25 ± 1.09	1.25 ± 1.09	0.50 ± 0.87	1.25 ± 0.83
<i>S. feltiae</i>	UNK-36	4.75 ± 0.43 ^a	3.50 ± 1.12 ^a	3.50 ± 1.12 ^a	2.50 ± 1.50 ^a	1.25 ± 1.09 ^a	1.00 ± 0.00 ^a	2.50 ± 1.12 ^a	1.75 ± 1.09 ^a
<i>S. feltiae</i>	SN	4.50 ± 0.87 ^a	3.25 ± 2.05 ^a	2.50 ± 2.50 ^a	3.75 ± 0.83 ^a	2.25 ± 2.28 ^a	1.00 ± 0.43 ^a	1.50 ± 1.12	1.00 ± 0.00
Control	—	0.25 ± 0.43	0.50 ± 0.87	0.50 ± 0.87	0.50 ± 0.50	0.25 ± 0.43	0.25 ± 0.43	0.25 ± 0.43	1.25 ± 0.83

Five wax moth larvae per treatment.

^a Nematode presence confirmed in ≥1 greater wax moth larva.

(1987a), who found *S. carpocapsae* (= *S. feltiae* Breton and *S. feltiae* Mexican) to be more virulent than *S. feltiae* (= *S. bibionis* KL and *S. bibionis* SN). Both our study and that of Mullens et al. (1987) found *Heterorhabditis* spp. to have low virulence, whereas Geden et al. (1986) reported higher virulence for *H. bacteriophora* (= *H. heliothidis*) than *S. carpocapsae* (= *S. feltiae* DD-136) and *S. glaseri*. *S. glaseri* had very low virulence in both Geden et al. (1986) and our study. Of the 8 *Heterorhabditis* species we tested, only 2—*H. bacteriophora* and *H. zealandica*—had been examined for virulence toward house flies in previous studies. Three of the 5 *Steinernema* species had been previously examined.

Serial passage through maggots for 10 generations did not change the pathogenicity of any of the *Steinernema* strains (Table 2). Although initially disappointing, these results indicate that nematode strains can be screened effectively for pathogenicity toward maggots, and possibly other hosts, without acclimation to that host for several generations. In addition, greater wax moth larvae can be used as hosts for *in vivo* mass production without reducing pathogenicity toward maggots.

Other studies have documented the pathogenicity of several strains of entomopathogenic nematodes toward house fly larvae in filter paper assays (Renn et al. 1985, Geden et al. 1986, Belton et al. 1987, Mullens et al. 1987a). However, all of those studies focused upon the use of these nematodes for the control of flies in chicken manure. The eventual failure of these programs was attributed to the nematodes inability to survive in chicken manure (Georgis et al. 1987). In contrast, we found that the nematodes not only survive and remain infective in manure for >1 mo, but that their pathogenicity is equal to, or greater than, that observed in filter paper assays (Tables 2 and 3). Our results appear to differ from those of Shapiro et al. (1996), who found manure reduced nematode virulence when added to soil as a fertilizer. However, Shapiro et al. examined only *S. carpocapsae*, a species we did not test in the manure substrate because it did not perform as well as *S. feltiae* in the filter paper assays. Bednarek and Gaugler (1997) found *S. feltiae* populations actually increased with the addition of manure to soils supporting our results.

Our persistence studies (Table 4) were conducted in the absence of hosts for nematode reproduction. Presumably, in a feedlot environment, fresh hosts will

be available for the nematodes to cycle continuously, thus not only extending the persistence of the nematodes, but possibly increasing the population with time. Based upon the LC₉₉ value of 104 nematodes per square-centimeter of surface area, we project an application rate of 1 million nematodes per square-meter of feedlot. We estimate (unpublished data) that ≈10,000 infective juvenile nematodes are produced per maggot. Hence, successful reproduction in 100 maggots per square-meter will maintain a nematode population adequate for 99% control. Skoda et al. (1996) estimated an average population of about 2,200 stable flies, *Stomoxys calcitrans* (L.), and 4,000 house flies or 6,200 flies total per square-meter throughout the feedlot. Therefore, only .15% of the infected maggots would need to support complete nematode development to maintain a constant population at the LC₉₉ level.

Acknowledgments

We thank R. J. Akhursts, M. A. Brooks, A. M. Burnell, H. E. Cabanillas, C. T. Griffin, J. J. Jackson, S. A. Joyce, J. Liu, P. H. Smits, S. P. Stock, and P. R. Westerman for providing nematode stocks. K. Pruess and B. Siegfried (University of Nebraska) provided helpful suggestions and critical reviews of the manuscript. This work was done in cooperation with the Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE, and is published as Journal Series, Nebraska Agricultural Research Division Paper No. 11924.

References Cited

- Bednarek, A. J., and R. Gaugler. 1997. Compatibility of soil amendments with entomopathogenic nematodes. *J. Nematol.* 29: 220-227.
- Belton, P., T. A. Rutherford, D. B. Trotter, and J. M. Webster. 1987. *Heterorhabditis heliothidis*: potential biological control agent of house flies in caged-layer poultry barns. *J. Nematol.* 19: 263-266.
- Dutky, S. R., J. V. Thompson, and G. E. Cantwell. 1964. A technique for the mass propagation of the DD-136 nematode. *J. Insect Pathol.* 6: 417-422.
- Fuxa, J. R. 1987. *Spodoptera frugiperda* susceptibility to nuclear polyhedrosis virus isolates with reference to insect migration. *Environ. Entomol.* 16: 218-223.
- Gaugler, R. 1988. Ecological considerations in the biological control of soil-inhabiting insects with entomopathogenic nematodes. *Agric. Ecosyst. Environ.* 24: 351-360.
- Geden, C. J., R. C. Axtell, and W. M. Brooks. 1986. Susceptibility of the house fly, *Musca domestica* (Diptera: Mus-

- cidae), to the entomogenous nematodes *Steinernema feltiae*, *S. glaseri* (Steinernematidae). *J. Med. Entomol.* 23: 326-332.
- Georgis, R., B. A. Mullens, and J. A. Meyer.** 1987. Survival and movement of insect parasitic nematodes in poultry manure and their infectivity against *Musca domestica*. *J. Nematol.* 19: 292-295.
- Klein, M. G.** 1990. Efficacy against soil-inhabiting insect pests, pp. 195-214. *In* R. Gaugler and H. K. Kaya [eds.], *Entomopathogenic nematodes in biological control*. CRC, Boca Raton, FL.
- Laumont, D., H. Mauleon, and A. Kermarrec.** 1979. Données nouvelles sur le spectre d'hotes et le parasitisme du nematode entomophage *Neoplectana carpocapsae*. *Entomophaga* 24: 113-127.
- Mullens, B. A., J. A. Meyer, and T. L. Cyr.** 1987a. Infectivity of insect-parasitic nematodes (Rhabditida: Steinernematidae, Heterorhabditidae) for larvae of some manure-breeding flies (Diptera: Muscidae). *Environ. Entomol.* 16: 769-773.
- Mullens, B. A., J. A. Meyer, and R. Georgis.** 1987b. Field tests of insect-parasitic nematodes (Rhabditida: Steinernematidae, Heterorhabditidae) against larvae of manure-breeding flies (Diptera: Muscidae) on caged-layer poultry facilities. *J. Econ. Entomol.* 80: 438-442.
- Petersen, J. J.** 1993. Biotic agents to control house flies and stable flies, pp. 70-77. *In* G. D. Thomas and S. R. Skoda [eds.], *Rural flies in the urban environment?* North Central Regional Research Publication No. 335.
- Poinar, G. O., Jr.** 1979. *Nematodes for biological control of insects*. CRC, Boca Raton, FL.
- 1990.** Taxonomy and biology of Steinernematidae and Heterorhabditidae, pp. 23-61. *In* R. Gaugler and H. K. Kaya [eds.], *Entomopathogenic nematodes in biological control*. CRC, Boca Raton, FL.
- Renn, N., G. Barson, and P. N. Richardson.** 1985. Preliminary laboratory tests with two species of entomophilic nematodes for control of *Musca domestica* in intensive animal units. *Ann. Appl. Biol.* 106: 229-233.
- SAS Institute.** 1985. *Users manual, version 6.0*. SAS Institute, Cary, NC.
- Shapiro, D. I., G. L. Tylka, and L. C. Lewis.** 1996. Effects of fertilizers on virulence of *Steinernema carpocapsae*. *Appl. Soil Ecol.* 3: 27-34.
- Skoda, S. R., G. D. Thomas, and J. B. Campbell.** 1996. Comparison of core sampling and pupal traps for monitoring immature stable flies and house flies (Diptera: Muscidae) in beef feedlot pens. *J. Econ. Entomol.* 89: 428-434.
- White, G. F.** 1927. A method for obtaining infective nematode larvae from cultures. *Science (Wash. D.C.)* 66: 302-303.

Received for publication 5 February 1998; accepted 6 July 1998.
